



**University of  
Zurich**<sup>UZH</sup>

**Zurich Open Repository and  
Archive**

University of Zurich  
University Library  
Strickhofstrasse 39  
CH-8057 Zurich  
[www.zora.uzh.ch](http://www.zora.uzh.ch)

---

Year: 2015

---

## **Paradoxical response to furosemide in uromodulin-associated kidney disease**

Labriola, Laura ; Olinger, Eric ; Belge, Hendrica ; Pirson, Yves ; Dahan, Karin ; Devuyst, Olivier

**Abstract:** Mutations in the UMOD gene coding for uromodulin cause autosomal dominant tubulointerstitial kidney disease. Uromodulin is known to regulate transport processes in the thick ascending limb, but it remains unknown whether UMOD mutations are associated with functional tubular alterations in the early phase of the disease. The responses to furosemide and to a water deprivation test were compared in a 32-year-old female patient carrying the pathogenic UMOD mutation p.C217G and her unaffected 31-year-old sister. A single dose of furosemide induced an intense headache with exaggerated decrease in blood pressure ( $\Delta$ syst: 30 versus 20 mmHg;  $\Delta$ diast: 18 versus 5 mmHg) and body weight ( $\Delta$ 2.6 kg versus  $\Delta$ 0.9 kg over 3 h) in the proband versus unaffected sib. The diuretic response and the fall in urine osmolality were also more important and detected earlier in the affected sib. Water deprivation led to increased plasma osmolality and urine concentration in both siblings; however, the response to desmopressin was attenuated in the affected sib. These data reveal that mutations of uromodulin cause specific transport alterations, including exaggerated response to furosemide and a failure to maximally concentrate urine, in the early phase of the disease.

DOI: <https://doi.org/10.1093/ndt/gfu389>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-104187>

Journal Article

Accepted Version

Originally published at:

Labriola, Laura; Olinger, Eric; Belge, Hendrica; Pirson, Yves; Dahan, Karin; Devuyst, Olivier (2015). Paradoxical response to furosemide in uromodulin-associated kidney disease. *Nephrology, Dialysis, Transplantation*, 30(2):330-335.

DOI: <https://doi.org/10.1093/ndt/gfu389>

10 Paradoxical Response to Furosemide in Uromodulin-Associated  
11  
12 Kidney Disease  
13  
14  
15

16 Laura Labriola<sup>1\*</sup>, Eric Olinger<sup>2\*</sup>, Hendrica Belge<sup>2</sup>, Yves Pirson<sup>1</sup>, Karin Dahan<sup>3</sup>  
17  
18 and Olivier Devuyst<sup>1,2</sup>  
19  
20

21  
22 <sup>1</sup> UCL Nephrology, Brussels, Belgium;

23 <sup>2</sup> Institute of Physiology, University of Zurich, Zurich, Switzerland;

24  
25 <sup>3</sup> Institute of Pathology and Genetics, Gosselies, Belgium  
26  
27

28  
29 \*LL and EO contributed equally to this study.  
30  
31

32 Correspondence: to OD, Institute of Physiology, ZIHP, University of Zurich, Switzerland;  
33 E-mail: olivier.devuyst@uzh.ch  
34  
35

36 Runningtitle: Uromodulin and TAL function  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56

## ABSTRACT

Mutations in the UMOD gene coding for uromodulin cause autosomal dominant tubulointerstitial kidney disease. Uromodulin is known to regulate transport processes in the thick ascending limb, but it remains unknown whether UMOD mutations are associated with functional tubular alterations in the early phase of the disease.

The responses to furosemide and to a water deprivation test were compared in a 32-year-old female patient carrying the pathogenic UMOD mutation p.C217G and her unaffected 31-year-old sister. A single dose of furosemide induced an intense headache with exaggerated decrease in blood pressure ( $\Delta$  syst: 30 vs. 20mmHg;  $\Delta$  diast: 18 vs. 5mmHg) and body weight ( $\Delta$  2.6kg vs.  $\Delta$  0.9kg over 3h) in the proband vs. unaffected sib. The diuretic response and the fall in urine osmolality were also more important and detected earlier in the affected sib.

Water deprivation led to increased plasma osmolality and urine concentration in both siblings; however, the response to desmopressin was attenuated in the affected sib. These data reveal that mutations of uromodulin cause specific transport alterations, including exaggerated response to furosemide and a failure to maximally concentrate urine, in the early phase of the disease.

## INTRODUCTION

Uromodulin (Tamm-Horsfall protein) is exclusively produced in the epithelial cells lining the thick ascending limb (TAL) of Henle's loop and is the most abundant protein in the normal urine. Evidence obtained from in vitro studies and knock-out mouse studies have shown that uromodulin protects against urinary tract infection and kidney stones, and regulates transport systems operating in the TAL (1). The latter include the sodium-chloride cotransporter NKCC2 and the potassium channel ROMK, which operate in parallel and are essential to reabsorb 25% of the filtered NaCl and to mediate the urinary concentrating ability (2).

Mutations in the UMOD gene encoding uromodulin have been shown to cause medullary cystic kidney disease type 2 (MCKD2; MIM 603860), familial juvenile hyperuricemic nephropathy (FJHN; MIM 162000) and glomerulocystic kidney disease (GCKD; MIM 609886) (3-5) which are collectively referred to as uromodulin-associated kidney diseases (UAKD). UAKD are autosomal dominant disorders characterised by hyperuricemia and gout early in life, alteration of urinary concentrating ability, and tubulo-interstitial fibrosis with occasional cysts at the cortico-medullary junction (1). UAKD invariably lead to chronic renal failure during the third to seventh decade of life (6). There is no specific therapy to slow renal disease progression, and most patients end up with ESRD.

The fact that hyperuricemia is the most consistent feature observed in patients harbouring UMOD mutations has led to suggest that the disease causes a dysfunction of the TAL, with NaCl-loss and secondary reabsorption of uric acid in the proximal tubule (6,7). This hypothesis has been supported by functional defects evidenced in the first transgenic mouse model of disease (8). However, the existence of a specific dysfunction of the TAL as an early consequence of defective uromodulin processing has not been evaluated in patients harbouring a pathogenic UMOD mutation.

To investigate whether a pathogenic UMOD mutation is associated with functional alterations in the TAL in the early stage of the disease, we performed a family-based study to compare the response to a furosemide test and to a water-deprivation test with administration of desmopressin in a proband carrying a pathogenic mutation of UMOD and her unaffected sister.

## PATIENTS AND METHODS

The proband, a 32-year-old female harbouring a missense (p.C217G) mutation in the UMOD gene (III,2), and her 31-year-old sister tested negative for the mutation (III,4) belong to a four-generation family. The pathogenic nature of the p.C217G mutation of UMOD was evidenced by the typical course of UAKD observed in the other sister of the proband (III,3) who reached end-stage renal disease at age 37 years ([Fig.1A](#)). The proband and her unaffected sister had an eGFR (CKD-EPI)  $\geq 60$  ml/min/1.73 m<sup>2</sup> at time of testing, and none of them was diabetic. They were not taking any drug, including diuretics, allopurinol, or antihypertensive medications. The protocol was approved by the Ethical Committee of the UCL Medical School (Brussels), and informed consent was provided.

A 24-h urine collection was obtained in both sisters immediately before the furosemide test. The day of the furosemide test, a baseline clinical and biological evaluation was performed at 9:30 AM, 4h before giving a single oral dose of 80 mg furosemide. Baseline values for the furosemide test ([Fig.1B](#)) correspond to a 3-h urine collection preceding immediately the administration of furosemide. After furosemide administration, urine samples were collected hourly during 3 h and analysed for Na<sup>+</sup> and K<sup>+</sup>, creatinine, urea and uric acid concentrations, osmolality and pH. Clinical and biological evaluations were repeated at the end of the test (time +3h; see [Table1](#)). The following day (i.e. after 18h of normal hydration), both individuals underwent a 8-h water deprivation test, including a 30-min intravenous

1  
2  
3 infusion of desmopressin acetate (Minirin<sup>R</sup>, 0.3 µg/kg; Ferring AG, Baar, Switzerland) after 6  
4  
5 h of water deprivation. Desmopressin (or deamino-8D-arginine vasopressin, dDAVP) is a  
6  
7 peptidic analogue of endogenous antidiuretic hormone (ADH) with a higher affinity for V2  
8  
9 receptors and negligible vasopressive potential. A clinical evaluation was performed hourly  
10  
11 and plasma and urine samples were taken every 2 h during the water deprivation and 60 and  
12  
13 120 min after initiation of desmopressin perfusion. Dietary intake was identical in the two  
14  
15 sisters before the furosemide test and throughout the water deprivation test.  
16  
17  
18  
19

## 20 21 RESULTS

22  
23  
24 The baseline osmolar balance in the two sisters immediately before the functional tests  
25  
26 was similar, thus allowing direct interindividual comparison of the responses to furosemide  
27  
28 and to water restriction (Values for affected sib vs. control sib: 24h Na<sup>+</sup> excretion: 228 mmol  
29  
30 vs. 307 mmol; 24h urea excretion: 29.1g vs. 23.1g; osmolar excretion rate: 779.2 µosm/min  
31  
32 vs. 787.5 µosm/min; plasma renin activity: 1.1 ng/mL/h vs. 0.8 ng/mL/h; plasma aldosterone:  
33  
34 0.31 nM vs. 0.30 nM, respectively. Plasma ADH: <0.2 pg/mL in both sibs).  
35  
36

37  
38 The furosemide test was poorly tolerated by the proband, who reported intense  
39  
40 headache, a severe drop in blood pressure (Δ30/Δ18 mmHg vs. Δ20/Δ5 mmHg in the  
41  
42 unaffected sister) and a weight loss of 2.6 kg (vs. weight loss of 0.9 kg in the unaffected  
43  
44 sister) in 3 h (Table1). After furosemide administration, both siblings showed the expected  
45  
46 increase in diuresis with concomitant fall in urine osmolality. However, the changes were  
47  
48 more important and were detected earlier in the proband, who also showed a more  
49  
50 pronounced and earlier increase of fractional excretion of Na<sup>+</sup> than the control sibling (Fig.  
51  
52 1B).  
53  
54

55  
56 The water deprivation test was well tolerated by both participants. Blood pressure and  
57  
58 weight remained constant during all the procedure. As expected, water deprivation resulted in  
59  
60

1  
2  
3 a progressive increase in plasma osmolality with an increased urine osmolality and a reduced  
4  
5 urine flow in both subjects (Fig.1B). The response to desmopressin was normal in the control  
6  
7 sibling, with an increase in urine osmolality (from 983 to 1124 mOsm/kg H<sub>2</sub>O, 2h after  
8  
9 infusion), a decrease in urine flow (from 30 to 18 mL/h), and a decrease in plasma osmolality  
10  
11 (from 298 to 295 mOsm/kg H<sub>2</sub>O). In contrast, the affected sib showed a blunted increase in  
12  
13 urine osmolality (from 834 to 869 mOsm/kg H<sub>2</sub>O), an increase in urine flow (from 17.5 mL/h  
14  
15 to 20 mL/h), and a paradoxical increase in plasma osmolality (from 298 to 305 mOsm/kg  
16  
17 H<sub>2</sub>O) (Fig.1B).  
18  
19

20 Pathology examination of the end-stage kidney biopsy of patient (III,3) (Fig.1C)  
21  
22 revealed marked tubulo-interstitial lesions, with thickening of the tubular basement membrane  
23  
24 and abnormal processing and accumulation of uromodulin in the tubular cells. Of note, there  
25  
26 was no detectable immunostaining for NKCC2 in the biopsy.  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

## DISCUSSION

In this study, we performed a sib-pair functional testing to evaluate whether pathogenic mutations of UMOD are associated with dysfunction of the TAL in the early phase of UAKD. Our data reveal that the proband carrying a UMOD mutation shows a clinically and biologically exaggerated response to furosemide and a failure to maximally concentrate urine after desmopressin administration. To the best of our knowledge, this is the first report of abnormal tubular functional testing in a subject harbouring a pathogenic UMOD mutation.

Several lines of evidence suggest that uromodulin plays an important role in regulating NKCC2 and ROMK in the TAL (9-11). These two transport processes mediate the furosemide-sensitive NaCl reabsorption and the generation of the osmotic gradient which drives the urine concentrating ability. Patients with UMOD mutations are characterized by an early defect in urine concentration (3-5), which could be paralleled with a discrete NaCl-losing phenotype explaining hyperuricemia (6,7). Studies of mice harbouring a pathogenic mutation of uromodulin revealed that the urinary concentrating defect precedes renal failure and is related to a specific defect in the TAL (8). The latter includes defective expression of NKCC2 and other markers, secondary to the accumulation of mutant uromodulin in the endoplasmic reticulum (ER) (8). We confirm such lesions in the end-stage kidney biopsy of patient III,3, showing extensive tubulo-interstitial damage with abnormal processing of uromodulin and loss of NKCC2 immunoreactivity.

The demonstration of an exaggerated response to furosemide in the proband, despite the lower eGFR than her control sister, is of particular interest. Since uromodulin expression levels regulate NKCC2 activity (10, 11), it is conceivable that the reduced trafficking of native uromodulin in UAKD might lead to a decreased amount of active NKCC2 on the apical cell membrane. This effect would be amplified by the fact that disease-causing UMOD mutations also lead to decreased surface expression of ROMK (9). The paradoxically



1  
2  
3 increased response to furosemide, as observed here, might therefore indicate that there is still  
4  
5 sufficient NaCl uptake capacity (e.g. due to hyperactivated residual NKCC2) to maintain a  
6  
7 steady-state in the early stage of disease. Administration of furosemide may disrupt this  
8  
9 equilibrium, by inhibiting active NKCC2 transporters with no immediately available  
10  
11 compensation in other tubule segments. The normal serum uric acid levels (Table1) and the  
12  
13 slightly higher uric acid fractional excretion in the proband (5.9% vs. 5.1% in the healthy sib  
14  
15 calculated from the baseline 3-h urine collection) may indeed suggest a maintained sodium  
16  
17 balance by residual TAL function without proximal compensatory adaptation.  
18  
19

20  
21 Alternatively, the exaggerated response to furosemide in the proband with early  
22  
23 UAKD could unmask a shift of steady- state NaCl reabsorption inside the TAL. Based on  
24  
25 functional and anatomical differences, the TAL segment could indeed be sub-divided into a  
26  
27 cortical (cTAL) and a medullary (mTAL) portion. The cTAL, which lies in an environment  
28  
29 isosmotic to the plasma, is responsible for the majority of NaCl reabsorption and is the true  
30  
31 “diluting segment” of the nephron (12, 13). In the rat kidney, two different cell types coexist  
32  
33 in the TAL: cells with a smooth surface and a dense subapical NKCC2-containing vesicle  
34  
35 pool, which predominate in the mTAL, and cells with a rough surface and a much less  
36  
37 abundant vesicle system, which largely dominate in the cTAL (14). Although human UMOD  
38  
39 transcript levels have been found quite similar in mTAL and cTAL (15), human uromodulin  
40  
41 protein levels were 4-fold greater in the TAL-enriched outer medulla than in the cortex by  
42  
43 immunoblotting (16). In healthy individuals, steady state NaCl reabsorption is probably  
44  
45 achieved by the early part of cTAL, leaving a “physiological reserve” in the late part of the  
46  
47 cTAL (12). In the proband with early UAKD, a reduced function of NKCC2 in mTAL could  
48  
49 be compensated by a larger implication of the cTAL, possibly spared from the transport  
50  
51 alterations associated with UAKD. Thus, a shift of NaCl reabsorption to downstream/late part  
52  
53 of the cTAL may have used up the “physiological reserve” in the affected sib, which could  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 then contribute to the differential response to furosemide. The fact that, in contrast to the  
4  
5 control sib, the proband was unable to lower its urinary sodium concentration below plasma  
6  
7 sodium (data not shown) might further indicate that the diluting function of the TAL was  
8  
9 completely abolished by furosemide. Taken together, these data indicate that the response to  
10  
11 furosemide in UAKD patients is diphasic, with an exaggerated response in early disease and  
12  
13 an anticipated loss of response as predicted by the loss of NKCC2 in end-stage kidney. Loop  
14  
15 diuretics in these patients should thus be managed with great caution.  
16  
17

18  
19 At baseline, the proband showed a trend for increased plasma osmolality contrasting  
20  
21 with lower urine osmolality, compatible with a slight alteration of water homeostasis. The  
22  
23 observation of a preserved response to water deprivation followed by a blunted response to  
24  
25 desmopressin may again indicate a reduced “physiological reserve” of the urinary  
26  
27 concentrating ability in early UAKD. The normal response of the proband to water  
28  
29 deprivation is distinct from the severe nephrogenic diabetes insipidus observed in the  
30  
31 transgenic mutant mouse model (8). The difference is probably due to the fact that the tubular  
32  
33 and interstitial lesions observed in the patient are less severe than in the transgenic mice  
34  
35 analysed at a late stage of the disease. Additionally the water restriction test challenges mostly  
36  
37 the inner medulla with its ADH-sensitive urea transporters in the terminal collecting duct,  
38  
39 where uromodulin expression is absent (13). Possible explanations for the blunted response to  
40  
41 desmopressin include (i) a failure to increase or maintain interstitial medullary  
42  
43 hyperosmolality; and/or (ii) a decreased response to desmopressin. A less active osmolar  
44  
45 gradient generator, namely NKCC2 transporter, or a leaky TAL allowing water influx into the  
46  
47 interstitium or NaCl backleak into the lumen could underlie the former hypothesis.  
48  
49

50  
51 Considering the interactions of UMOD with NKCC2 and the physical properties (including  
52  
53 water impermeability) of uromodulin polymers formed under specific ionic conditions (1), an  
54  
55 altered interstitial osmolality could be very well explained by a reduced excretion of  
56  
57  
58  
59  
60

1  
2  
3 uromodulin. In fact, the Umod KO mice showed no increase in pNKCC2 after stimulation  
4  
5 with dDAVP, suggesting that uromodulin is important for the TAL sensitivity to vasopressin  
6  
7 (10).  
8

9  
10 The main limitations of this study include the small number of siblings involved and  
11  
12 the lack of histopathology data in the early stage of disease. However, we feel that these  
13  
14 observations give insights into the biology of uromodulin and the pathophysiology of UAKD.  
15  
16 We investigated a patient with preclinical disease, with presumably limited interstitial  
17  
18 inflammation/fibrosis (normal urinary sediment), and an unaffected sibling of the same  
19  
20 gender and similar age. The deleterious effect of the mutation was evidenced by the end-stage  
21  
22 kidney biopsy of a third sibling.  
23

24  
25 In conclusion, our study suggests that, in the early phase of UAKD, the lack of  
26  
27 functional uromodulin leads to a discrete dysfunction of the TAL, with maintenance of a  
28  
29 precarious clinical equilibrium that can be disturbed by specific testing. As the disease  
30  
31 progresses, the aggravation of the tubulo-interstitial lesions leads to overt tubular dysfunction  
32  
33 and compensatory mechanisms, culminating with chronic kidney disease.  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

## ACKNOWLEDGEMENTS

We acknowledge Mrs. Y. Cnops, Prof. J-P. Cosyns and Mrs. N. Van Oost for their help, and the reviewers for their helpful suggestions and comments. These studies were supported by the European Community's 7th Framework Programme (FP7/2007-2013) under grant agreement n° 246539 and 608847 (IKPP Marie Curie) and grant n° 305608 (EUREnOmics); Action de Recherche Concertée (ARC10/15-029, Communauté Française de Belgique); the FNRS and FRSM; Inter-University Attraction Pole (IUAP, Belgium Federal Government); supported by the Fonds National de la Recherche, Luxembourg (6903109); the NCCR Kidney.CH program (Swiss National Science Foundation); the Gebert Rief Stiftung (Project GRS-038/12); and the Swiss National Science Foundation 310030-146490.

Competing financial interests: The authors declare no competing financial interests.

## REFERENCES

1. Rampoldi L, Scolari F, Amoroso A, Ghiggeri GM, Devuyst O. The rediscovery of uromodulin (Tamm-Horsfall protein): from tubulointerstitial nephropathy to chronic kidney disease. *Kidney Int* 2011; 80: 338-347
2. Devuyst O. Salt wasting and blood pressure. *Nat Genet* 2008; 40: 495-496
3. Hart TC, Gorry MC, Hart PS et al. Mutations of the UMOD gene are responsible for medullary cystic kidney disease 2. *J Med Genet* 2002; 39: 882-892
4. Dahan K, Devuyst O, Smaers M et al. A cluster of mutations in the UMOD gene causes Familial Juvenile Hyperuricemic Nephropathy with abnormal expression of uromodulin. *J Am Soc Nephrol* 2003; 14: 2883-2893
5. Rampoldi L, Caridi G, Santon D, et al. Allelism of MCKD, FJHN and GCKD caused by impairment of uromodulin export dynamics. *Hum Mol Genet* 2003; 12: 3369-3384
6. Bollée G, Dahan K, Flamant M, et al. Phenotype and outcome in hereditary tubulointerstitial nephritis secondary to UMOD mutations. *Clin J Am Soc Nephrol* 2011; 6: 2429-2438
7. Scolari F, Caridi G, Rampoldi L et al.: Uromodulin Storage Diseases: Clinical Aspects and Mechanisms. *Am J Kidney Dis* 2004; 44: 987-999
8. Bernascone I, Janas S, Ikehata M, et al. A transgenic mouse model for uromodulin-associated kidney diseases shows specific tubulo-interstitial damage, urinary concentrating defect and renal failure. *Hum Mol Genet* 2010; 19: 2998-3010
9. Renigunta A, Renigunta V, Saritas T, Decher N, Mutig K, Waldegger S. Tamm-Horsfall glycoprotein interacts with renal outer medullary potassium channel ROMK2 and regulates its function. *J Biol Chem* 2010; 286: 2224-2235
10. Mutig K, Kahl T, Saritas T, et al. Activation of the bumetanide-sensitive Na<sup>+</sup>,K<sup>+</sup>,2Cl<sup>-</sup>-cotransporter (NKCC2) is facilitated by Tamm-Horsfall protein in a chloride-sensitive manner. *J Biol Chem* 2011; 286: 30200-30210
11. Trudu M, Janas S, Lanzani C, et al. Common noncoding UMOD gene variants induce salt-sensitive hypertension and kidney damage by increasing uromodulin expression. *Nat Med* 2013; 19: 1655-1660
12. Burg M. Thick ascending limb of Henle's loop. *Kidney Int* 1982; 22: 454-464
13. Bankir L, De Rouffignac C. Urinary concentrating ability : insights from comparative anatomy. *Am J Physiol* 1985; 249: 643-666
14. Nielsen S, Maunsbach AB, Ecelbarger CA, Knepper MA. Ultrastructural localization of Na-K-2Cl cotransporter in thick ascending limb and macula densa of rat kidney. *Am J Physiol* 1998; 275: 885-893

1  
2  
3 15. Chabardes-Garonne D, Mejean A, Aude JC, et al. A panoramic view of gene expression in the  
4 human kidney. Proc Natl Acad Sci USA 2003; 100: 13710-13715  
5

6 16. Serafini-Cessi F, Malagolini N, Cavallone D. Tamm-Horsfall glycoprotein: biology and clinical  
7 relevance. Am J Kidney Dis 2003; 42: 658-676  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

Table 1. Clinical and biological response to furosemide in the proband and the unaffected sibling.

The parameters were similarly recorded 4 h before (Baseline) and 3 h after furosemide administration (Furosemide +3h).

	<u>Proband(III.2)</u>		<u>UnaffectedSib(III.4)</u>	
	Baseline	Furosemide +3h	Baseline	Furosemide +3h
Weight (kg)	76.1	73.5	81.1	80.2
Blood pressure (mmHg)	140/90	110/72	135/80	115/75
eGFR (CKD-EPI; mL/min/1.73m <sup>2</sup> )	60	60	99	99
Plasma Urea (mg/dL) Plasma	45	43	25	23
Sodium (mmol/L) Plasma	141	138	139	138
Potassium (mmol/L) Plasma	4.4	3.7	3.8	3.8
Uric Acid (mg/dL) Plasma	6.3	6.3	6.4	6.4
Osmolality (mOsm/L) Plasma	298	298	287	288
Bicarbonate (mmol/L)	27.5	29.5	22.5	27.5

## Figure Legends

Figure 1. Pedigree of the family, UMOD mutation, response to furosemide and water deprivation and end-stage kidney biopsy.

A. Pedigree of the family and UMOD mutation. Individuals with a history of renal disease are depicted by black symbols. Circles denote females, squares males. The proband carrying the UMOD mutation (III,2) and her unaffected sister (III,4) are marked by an arrow and an arrowhead, respectively. Individuals tested for mutation are tagged by an asterisk (\*). Patient III,3 showed a typical course of uromodulin-associated kidney disease, with hyperuricemia at age 21 years and end-stage renal disease at age 37 years. Sequence analysis of the UMOD gene revealed a thymine to guanine transition at nucleotide 794 resulting in the substitution of a well conserved cysteine by a glycine at position 217 (p.C217G). Encoded amino acid sequence is indicated above the DNA sequence. Mutated nucleotide is boxed. The mutation has been previously reported by Dahan et al. (4).



B. Biological parameters at baseline and evaluation after furosemide administration and water deprivation with desmopressin perfusion. The patient harbouring the UMOD mutation is indicated by red dots and her unaffected sister by blue squares. Baseline values in the furosemide test have been obtained from a 3-hour urine collection immediately preceding furosemide administration. This same baseline diuresis has been plotted in the water deprivation test curve. Urine and plasma osmolality values at time 0 of the water deprivation test provide from urine and blood sample collected before the test.

C. End-stage kidney biopsy from patient (III,3) harbouring the p.C217G UMOD mutation. Hematoxylin-eosine staining (panel a) reveals interstitial fibrosis and tubular atrophy with abundant inflammatory infiltrate and the characteristic thickening of the tubular basement membrane (inset, arrows). Immunostaining (panel b) reveals diffuse intracellular accumulation of uromodulin in a subset of cells lining enlarged tubules (arrow) as well as cells displaying the normal apical staining for uromodulin (arrowhead). There is no detectable immunostaining for NKCC2 (panel c) in the kidney of patient (III,3) as compared with normal staining pattern (inset).

Original magnifications: a, x350; b, x200, c, x700.



Figure – panel A



Unaffected  
(III,4)




25  
26  
27  
28  
29

Affected  
(III,2)

Figure – panel A  
Figure 1- panel B

● Affected Sib (p.C217G)  
■ Control Sib

Furosemide Test



Water Deprivation Test

310 29 305  
30 30  
31 300

32 35  
33 36 295 290  
34 37

38  
39 285  
40

41  
42

